



For all you seq...
.




RNA Transcription



Methylation




DNA-Protein Interactions



- | References | |
|---|--|
| 3-C | Duan Z. et al. (2012) A genome-wide 3C-method for characterizing the three-dimensional architectures of genomes. <i>Methods</i> 58: 277-288 |
| 4-C | Zhao Z. et al. (2006) Circular chromosome conformation capture (4C) uncovers extensive networks of epigenetically-regulated intra- and interchromosomal interactions. <i>Nat Genet</i> 38: 1341-1347 |
| 5-C | Dostie J. et al. (2007) Mapping networks of physical interactions between genomic elements using 5C technology. <i>Nat Protoc</i> 2: 988-1002 |
| Bs-Seq | Meissner A. et al. (2008) Genome-scale DNA methylation maps of pluripotent and differentiated cells. <i>Nature</i> 454: 766-770 |
| Cap-Seq | Illingworth R. S. et al. (2010) Orphan CpG islands identify numerous conserved promoters in the mammalian genome. <i>PLoS Genet</i> 6: e1001134 |
| Chia-PET | Li G. et al. (2010) ChIA-PET tool for comprehensive chromatin interaction analysis with paired-end tag sequencing. <i>Genome Biol</i> 11: R22 |
| ChIP-Seq | Barski A. et al. (2007) High-resolution profiling of histone methylations in the human genome. <i>Nat Methods</i> 7: 995-1001 |
| ChIRP-Seq | Chu C. et al. (2011) Genomic maps of long noncoding RNA occupancy reveal principles of RNA-chromatin interactions. <i>Mol Cell</i> 44: 667-678 |
| CIP-TAP | Gu W. et al. (2012) CapSeq and CIP-TAP identify Pol II start sites and reveal capped small RNAs as C. elegans piRNA precursors. <i>Cell</i> 151: 1488-1500 |
| CLASH | Kudla G. et al. (2011) Cross-linking, ligation, and sequencing of hybrids reveals RNA-RNA interactions in yeast. <i>Proc Natl Acad Sci U S A</i> 108: 10010-10015 |
| Digital RNA | Shiroguchi K. et al. (2012) Digital RNA sequencing minimizes sequence-dependent bias and amplification noise with optimized single-molecule barcodes. <i>Proc Natl Acad Sci U S A</i> 109: 1347-1352 |
| DNase-Seq | McDaniell R. et al. (2010) Heritable individual-specific and allele-specific chromatin signatures in humans. <i>Science</i> 328: 235-239 |
| FAIRE-Seq | Gaulton K. J. et al. (2010) A map of open chromatin in human pancreatic islets. <i>Nat Genet</i> 42: 255-259 |
| Frag-Seq | Underwood J. G. et al. (2010) FragSeq: transcriptome-wide RNA structure probing using high-throughput sequencing. <i>Nat Methods</i> 7: 995-1001 |
| OXBS-Seq | Zong C. et al. (2012) Genome-wide detection of single-nucleotide and 5-hydroxymethylcytosine at single-base resolution. <i>Science</i> 336: 934-937 |
| GMUCT | Booth M. J. et al. (2011) Multiplexed RNA structure characterization with selective 2'-hydroxyl copy-number variations of a single human cell. <i>Science</i> 336: 11063-11068 |
| PAR-CLIP | Ponts N. et al. (2008) Identification of microRNAs and other small regulatory RNAs using cDNA library sequencing. <i>Methods</i> 44: 3-12 |
| smMIP | Hafner M. et al. (2013) Single molecule molecular inversion probes for targeted, high-accuracy detection of low-frequency variation. <i>Genome Res</i> 23: 843-854. |
| PARE-Seq | Hiatt JB (2013) Comprehensive mapping of long-range interactions reveals folding principles of the human genome. <i>Science</i> 326: 289-293 |
| Hi-C | Lieberman-Aiden E. et al. (2009) Global identification of microRNA-target RNA pairs by parallel analysis of RNA ends. <i>Nat Biotechnol</i> 26: 941-946 |
| PARS | Wan Y. et al. (2013) Genome-wide mapping of RNA structure using nuclease digestion and high-throughput sequencing. <i>Nat Protoc</i> 8: 849-869 |
| Histone Methylation | Down T. A. et al. (2008) A Bayesian deconvolution strategy for immunoprecipitation-based DNA methylome analysis. <i>Nat Biotechnol</i> 26: 779-785 |
| RC-Seq | Barbski A. et al. (2007) High-resolution profiling of histone methylations in the human genome. <i>Cell</i> 129: 823-837 |
| ICE | Baillie J. K. et al. (2011) Somatic retrotransposition alters the genetic landscape of the human brain. <i>Nature</i> 479: 534-537 |
| Ribo-Seq | Sakurai M. et al. (2010) Inosine cyanoethylation identifies A-to-I RNA editing sites in the human transcriptome. <i>Nat Chem Biol</i> 6: 733-740 |
| iCLIP | Meyer K. D. et al. (2012) Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons. <i>Cell</i> 149: 1635-1646 |
| TRAP-Seq | Ingolia N. T. et al. (2009) Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. <i>Science</i> 324: 218-223 |
| MethylCap-Seq | Konig J. et al. (2010) iCLIP reveals the function of hnRNP particles in splicing at individual nucleotide resolution. <i>Nat Struct Mol Biol</i> 17: 909-915 |
| NET-Seq | Zhao J. et al. (2010) Genome-wide identification of polycomb-associated RNAs by RIP-seq. <i>Mol Cell</i> 40: 939-953 |
| UMI Method | Churchman L. S. et al. (2011) Nascent transcript sequencing visualizes transcription at nucleotide resolution. <i>Nature</i> 469: 368-373 |
| CLIP-Seq | van Opijnen T. et al. (2013) Transposon insertion sequencing: a new tool for systems-level analysis of microorganisms. <i>Nat Rev Microbiol</i> 11: 435-442 |
| HITS-CLIP | Mylykangas S. et al. (2011) Efficient targeted resequencing of human germline and cancer genomes by oligonucleotide-selective sequencing. <i>Nat Biotechnol</i> 29: 1024-1027 |
| MDA | Dean F. B. et al. (2001) Rapid amplification of plasmid and phage DNA using Phi 29 DNA polymerase and multiply-primed rolling circle amplification. <i>Genome Res</i> 11: 1095-1099 |
| Did we miss anything? Please contact DL - Publications@illumina.com | |

identifiers. Nat Methods 9: 72-74



Sequencing by Synthesis

